


Stoichiometry: amounts of substances in balanced chemical reactions



How many moles of CO_2 can be produced from the combustion of 8.7 mol octane (C_8H_{18})?
(Use coefficients from balanced chemical equation to make a mole ratio)

How many grams of CO_2 can be produced from the combustion of 4.50×10^4 g of octane?

Limiting reactants

Whenever amounts of more than one reactant are known, you must find the limiting reactant

- Which reactant is consumed first?
- Calculate moles of a single product that each reactant will make
- Reactant that makes fewer product moles is the limiting reactant.
- That's how many product moles can be formed.



If 0.30 mol Zn react with 0.52 mol HCl, how many mol H₂ can be formed?

To calculate amount of leftover reactant, first calculate amount of excess reactant that was actually used, then subtract the given amount.

Limiting reactant with starting masses

If 7.36 g Zn react with 6.45 g S₈, how many grams ZnS can be produced? What mass of reactant remains?



Theoretical yield: calculated product mass from stoichiometry calculation

Actual yield: measured product mass from experiment

$$\% \text{ Yield} = \frac{\text{actual}}{\text{theoretical}} \times 100\%$$

Solution concentration and stoichiometry

Most chemical reactions in this course take place in solution (dissolved in water)


$$\text{Any concentration} = \frac{\text{amount solute}}{\text{amount solution}}$$

$$\text{Molarity (M)} = \frac{\text{moles of solute}}{\text{liters of solution}}$$

For instance, a 0.50 M $\text{AgNO}_3(aq)$ solution is called a 0.50 "molar" silver nitrate solution

1 liter of this solution contains _____ moles of AgNO_3

How do you prepare 250.0 mL of a 0.10 M $\text{AgNO}_3(aq)$ solution? (Use the given molarity as a conversion factor between moles solute and liters solution)



volumetric
flask

Concentration calculations

7.5 g CuCl_2 are dissolved in water to make 500.0 mL of solution. What is the molar concentration of this solution?

A reaction calls for 0.241 g K_2CO_3 . How many mL of 0.125 M $\text{K}_2\text{CO}_3(aq)$ should be added?


Dilution

Dilution: solvent is added to make a solution more dilute (less concentrated)

When adding water to a solution, what happens to the number of moles of solute?

Dilution equation: $M_1V_1 = M_2V_2$

Concentrated sulfuric acid is 18.0 M $\text{H}_2\text{SO}_4(aq)$. How do you make 10.0 L of 1.50 M $\text{H}_2\text{SO}_4(aq)$ by dilution?



Molarity converts between...

So, molarities can be used in stoichiometry problems along with the mole ratio from the balanced chemical equation.


How many grams of lead(II) iodide can be formed by mixing 1.0 mL of 0.50 M lead(II) nitrate solution with 2.0 mL 0.30 M sodium iodide solution?



Electrolytes and nonelectrolytes

Electrolyte: solute that causes solution to conduct electricity

Nonelectrolyte: solute that does not caused solution to conduct electricity



Solute

Electrolyte? Nonelectrolyte?

Deionized water

NaCl

$C_{12}H_{22}O_{11}$ (sugar)

NH_4Cl


C_3H_6O

Dissolving of electrolytes

Electrolytes **dissociate into ions** when dissolving




Dissolution of an Ionic Compound



Copyright © 2008 Pearson Prentice Hall, Inc.

Nonelectrolytes **remain as neutral molecules** when dissolving

Sugar Solution



Copyright © 2008 Pearson Prentice Hall, Inc.


Acids, Strong/weak electrolytes

Acids are molecular substances that dissociate when dissolved to release H^+ ions.

Strong electrolytes dissociate completely into ions:




Weak electrolytes do not completely dissociate into ions:



Solubility of ionic compounds

Not all ionic compounds dissolve in water!

- The ones that **do** are strong electrolytes
- The ones that **do not dissolve** remain solid in their ionic lattice when added to water



Solubility Rules for Ionic Compounds

The following table will be given on the exam **without** the formulas in parentheses.

Compounds Containing the Following Ions Are Mostly Soluble

Li^+ , Na^+ , K^+ , NH_4^+

nitrate (NO_3^-), acetate ($\text{C}_2\text{H}_3\text{O}_2^-$)

chloride (Cl^-), bromide (Br^-), iodide (I^-)

sulfate (SO_4^{2-})

Exceptions

None

None

When any of these ions pairs with Ag^+ , Hg_2^{2+} , or Pb^{2+} , the compound is insoluble

When sulfate pairs with Sr^{2+} , Ba^{2+} , Pb^{2+} , or Ca^{2+} the compound is insoluble

Compounds Containing the Following Ions Are Mostly Insoluble

hydroxide (OH^-), sulfide (S^{2-})

Exceptions

When either of these ions pairs with Li^+ , Na^+ , K^+ , or NH_4^+ , the compound is soluble

When sulfide (S^{2-}) pairs with Ca^{2+} , Sr^{2+} , or Ba^{2+} , the compound is soluble

When hydroxide (OH^-) pairs with Ca^{2+} , Sr^{2+} , or Ba^{2+} , the compound is slightly soluble (for many purposes, these may be considered insoluble)


carbonate (CO_3^{2-}), phosphate (PO_4^{3-})

When either of these ions pairs with Li^+ , Na^+ , K^+ , or NH_4^+ , the compound is soluble

Precipitation reactions

Precipitation reaction: two ionic compounds trade their ions to produce a solid, insoluble product


Write the balanced chemical equation with phase labels for reaction of solutions of potassium iodide and lead(II) nitrate.



If all products are soluble, **NO REACTION** occurs!

Molecular, complete ionic, and net ionic equations

Molecular equation: balanced chemical equation showing neutral formulas for all reactants and products.



Complete ionic equation: all strong electrolytes are written as separate ions with their own coefficients and phase labels. (Pure solids, liquids, gases, weak electrolytes, and nonelectrolytes written as molecules.)

Spectator ions do not participate in the chemical reaction (shown the same on both sides)

Net ionic equation: omits spectator ions - only shows the chemical change that has occurred

(What happens in the net ionic equation when it's considered "no reaction?" Try it for the combination of NaI and CaNO_3)

Acid-base reactions

Acid: produces H^+ ions in aqueous solutions

Base: produces OH^- ions in aqueous solutions

Strong acids dissociate completely (they are strong electrolytes) - there are 6 common ones:



Weak acids are other formulas that start with H such as:





Strong bases are the soluble or slightly soluble hydroxides: NaOH , LiOH , KOH , $\text{Ca}(\text{OH})_2$, $\text{Ba}(\text{OH})_2$

Acid-base neutralization reactions are the combination of an acid and a base to produce water and a salt

Write an equation for the acid-base neutralization reaction of nitric acid and calcium hydroxide

Acid-base titrations

Titration: controlled addition of one reactant (with known concentration) to another to determine an unknown concentration



Copyright © 2008 Pearson Prentice Hall, Inc.

Equivalence point: reaction is complete - equal moles of both reactants completely react with each other

Acid-base indicator: changes color to signal endpoint (phenolphthalein changes pink in excess OH^-)


Titration problem

25.0 mL of $\text{HCl}(aq)$ of unknown concentration were titrated with 12.45 mL of 0.100 M $\text{NaOH}(aq)$. What is the molarity of the $\text{HCl}(aq)$?



Gas-evolution reactions

Sulfides, sulfites, and carbonates form a gaseous product when reacting with an acid



Oxidation-reduction reactions

Oxidation-reduction (redox) reaction: electrons are transferred from one reactant to the other

Oxidation numbers: keep track of electrons in a reaction

Rules for Assigning Oxidation Numbers (must be memorized)

Rule	Applies to	Statement
1	Elements	The oxidation number of an atom in an element is zero
2	Monatomic ions	The oxidation number of an atom in a monatomic ion equals the charge on the ion.
3	Oxygen	The oxidation number of oxygen is -2 in most of its compounds. (An exception is O in H_2O_2 and other peroxides, where the oxidation number is -1 .)
4	Hydrogen	The oxidation number of hydrogen is $+1$ in most of its compounds. (The oxidation number of hydrogen is -1 in binary compounds with a metal, such as CaH_2 .)
5	Halogens	The oxidation number of fluorine is -1 in all of its compounds. Each of the other halogens (Cl, Br, I) has an oxidation number of -1 in binary compounds, except when the other element is another halogen above it in the periodic table or the other element is oxygen.
6	Compounds and ions	The sum of the oxidation numbers of the atoms in a compound is zero. The sum of the oxidation numbers of the atoms in a polyatomic ion equals the charge on the ion.

What is the oxidation number of sulfur in Na_2SO_4 ?

Oxidation-reduction reactions


Oxidation: loss of electrons (oxidation # increases)

Reduction: gain of electrons (oxidation # decreases)

Oxidizing agent: the reactant that is itself reduced

Reducing agent: the reactant that is itself oxidized

Identify the elements that are oxidized and reduced, and identify the oxidizing agent and reducing agent.



Combustion reaction: an oxidation reduction reaction with O_2 as a reactant.

Carbon-containing compounds undergo combustion to form CO_2 and H_2O

Write the equation for the combustion of C_6H_6

Metals undergo combustion to form metal oxides.