

## Chapter 6: Thermochemistry

**Thermochemistry:** study of the relationships between chemistry and energy

**Energy:** capacity to do work

**Work:** result of a force acting over a certain distance, one way to transfer energy

Types of energy:

- **Kinetic energy:** energy of motion
  - **Thermal energy:** energy associated with temperature, transferred by **heat**
- **Potential energy:** energy of position or composition
  - **Chemical energy:** energy associated with the composition of chemical compounds

**Law of conservation of energy:** energy cannot be created or destroyed, only transferred

- One object to another
- One type of energy to another

**Units of energy:**

- joule (J) =  $1 \text{ kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$  (SI unit)
- calorie (cal) = 4.184 J (heat 1 g H<sub>2</sub>O · 1 °C)
- Calorie (Cal) = 1 kcal = 1000 cal (food labels)
- kilowatt-hour (kWh) =  $3.60 \times 10^6 \text{ J}$

## System and surroundings

The transfer of energy can be considered in terms of a system and its surroundings

- **System:** a specific place or substance
- **Surroundings:** everything outside of the system

In thermochemistry, the system is usually a **chemical reaction**, and it can use heat and work to transfer energy to or from its surroundings:

- Combustion of natural gas in a furnace
- Combustion of gasoline in a car's engine
- Chemical cold pack


## First law of thermodynamics

**1st law:** the total energy of the universe is constant

- law of conservation of energy
- perpetual motion?

**Internal energy ( $E$ ):** sum of all forms of kinetic and potential energy in a certain system

Internal energy is a state function: it depends **only** on the makeup of the system itself at one point in time, not its history (it doesn't add up over time)



The **change** in any state function can be considered:

$$\Delta E = E_{\text{final}} - E_{\text{initial}}$$

## Work and heat

The change in a system's internal energy,  $\Delta E$ , depends on how much work ( $w$ ) was done and how much heat was transferred ( $q$ ):

$$\Delta E = q + w \qquad (\Delta E = E_{\text{final}} - E_{\text{initial}})$$

**Signs** are always considered from the system's perspective.

If energy flows **out of** the system,  $\Delta E$  is:

If energy flows **into** the system,  $\Delta E$  is:

If heat flows **out of** the system,  $q$  is:

If heat flows **into** the system,  $q$  is:

## Measuring heat:

Heat always transfers thermal energy from areas of \_\_\_\_\_ temperature to areas of \_\_\_\_\_ temperature

Heat capacity ( $C$ ): heat required to raise the temperature of an object by  $1^\circ\text{C}$

Units:

Specific heat capacity ( $C_s$  or  $s$ ): heat required to raise the temperature of **1 g of a substance** by  $1^\circ\text{C}$

Units:

Calculating heat transfers

For an object,  $q =$

For a certain mass of a substance,  $q =$

How much energy is required to heat 50.0 g water from 25.0 °C to 100.0 °C?

**TABLE 6.4** Specific Heat Capacities of Some Common Substances

Substance	Specific Heat Capacity, $C_s$ (J/g · °C)*
<b>Elements</b>	
Lead	0.128
Gold	0.128
Silver	0.235
Copper	0.385
Iron	0.449
Aluminum	0.903
<b>Compounds</b>	
Ethanol	2.42
Water	4.18
<b>Materials</b>	
Glass (Pyrex)	0.75
Granite	0.79
Sand	0.84

\*At 298 K.

Copyright © 2008 Pearson Prentice Hall, Inc.


How long will this take a 1000 W hotplate (assuming all of the heat is transferred to the water)? (1 W = 1 J/s)

Reaction work

What if the system is a **chemical reaction** that gives off or absorbs heat?

$$\Delta E_{\text{rxn}} = q + w$$

The most common work a chemical reaction does is expanding gases against an exterior pressure (pressure-volume work).



Copyright © 2008 Pearson Prentice Hall, Inc.

$$w = -P\Delta V$$


But usually we are mostly interested in the heat exchanged with the surroundings

If  $\Delta V = 0$ ,  $w = 0$  and  $\Delta E_{\text{rxn}} = q_V$  (constant volume)

If  $P$  is constant, we can use a different state function called enthalpy to only consider heat transferred.

## Constant-volume calorimetry

A **bomb calorimeter** is used to measure  $\Delta E_{\text{rxn}}$



$$\Delta E_{\text{rxn}} = q + w$$

$$w = -P\Delta V$$

If  $\Delta V = 0$ ,  $w = 0$  and  $\Delta E_{\text{rxn}} = q_V$  (constant volume)

- $q$  is a measurable energy transfer for a certain experimental process
- State functions like  $\Delta E$  are more general - they're given for a balanced chemical equation or per mole of a substance

$$q_{\text{cal}} = C_{\text{cal}} \Delta t \quad (C_{\text{cal}} \text{ is given})$$

$$q_{\text{rxn}} = -q_{\text{cal}}$$

$$\Delta E_{\text{rxn}} = q_{\text{rxn}} / n$$

## Constant-volume calorimetry

$$q_{\text{cal}} = C_{\text{cal}} \Delta t \quad (C_{\text{cal}} \text{ is given})$$

$$q_{\text{rxn}} = -q_{\text{cal}}$$

$$\Delta E_{\text{rxn}} = (q_{\text{rxn}}) / n$$

2.14 g  $\text{C}_5\text{H}_{12}$  is combusted in a bomb calorimeter and the temperature rises from 25.50 °C to 41.23 °C. The heat capacity of the calorimeter is 6.23 kJ/°C. What is  $\Delta E$  for the combustion of  $\text{C}_5\text{H}_{12}$  in kJ/mol?

$\Delta H$ , Change in enthalpy

It's more convenient to do a reaction open to the atmosphere, where pressure is constant.

The heat exchanged in a specific experiment is  $q_p$  and its related state function for a general process is  **$\Delta H$**  : change in enthalpy.

Since  $\Delta H$  only involves the heat given off or absorbed, it has the same sign conventions as  $q$ :

**Endothermic reaction**: heat is transferred from surroundings to system (inwards)

$\Delta H$  is:

**Exothermic reaction**: heat is transferred from system to surroundings (outwards)

$\Delta H$  is:




Copyright © 2008 Pearson Prentice Hall, Inc.

Thermochemical equations

$\Delta H_{\text{rxn}}$  is always given per a balanced chemical equation.

**Thermochemical equation:**

- balanced chemical equation
- thermochemical state function ( $\Delta H_{\text{rxn}}$ )
- molar interpretation (coefficients = # moles)




How much heat (in kJ) is given off when 20.0 g  $\text{H}_2$  react with excess oxygen?

## Constant-pressure calorimetry

$\Delta H_{\text{rxn}}$  can be measured using a constant-pressure calorimeter

The Coffee-Cup Calorimeter



We assume all of the heat lost by the reaction is gained by the solution (or vice-versa), and since there is a varying amount of solution, we use its specific heat capacity ( $C_s$  or  $s$ )

$$q_{\text{soln}} = (m_{\text{soln}})(C_{s,\text{soln}})(\Delta t)$$

$$q_{\text{rxn}} =$$


If the reaction is made only of dilute solutions, you can make these assumptions:

- $C_{s,\text{soln}} = C_{s,\text{water}} = 4.184 \text{ J/g}^\circ\text{C}$
- $m_{\text{solutions}} = m_{\text{water}}$
- $d_{\text{solutions}} = 1 \text{ g/mL}$

## Constant-pressure calorimetry problem

50.0 mL of 0.500 M  $\text{H}_2\text{SO}_4(aq)$  were added to 20.0 mL of 0.500 M NaOH in a coffee-cup calorimeter. The temperature rose from 25.00 °C to 27.20 °C. Assume  $C_s = 4.184 \text{ J/g}^\circ\text{C}$ . What is  $q_{\text{rxn}}$  in kJ?

What is  $\Delta H_{\text{rxn}}$  for the following balanced chemical equation?



Find the limiting reactant (LR):


What is  $n_{\text{LR}}$  for the reaction that was performed?

$$\Delta H_{\text{rxn}} = \text{coefficient}_{\text{LR}} \times \frac{q_{\text{rxn}}}{n_{\text{LR}}}$$

### $\Delta H$ and Hess's Law

Thermochemical state functions like  $\Delta H$  are given for a certain chemical equation, where coefficients = # mol

#### 1. Doubling coefficients doubles $\Delta H$ , etc.



#### 2. Reversing an equation changes the sign of $\Delta H$



#### 3. Adding equations adds their $\Delta H$ values

Find  $\Delta H_{\text{rxn}}$  for the following reaction:



Use the following reactions with known  $\Delta H$  values:



### Hess's law practice

Find  $\Delta H_{\text{rxn}}$  for the following reaction:



Use the following reactions with known  $\Delta H$  values:



Standard enthalpies of formation

**Standard thermodynamic states:**

- gases: 1 atm
- liquids or solids: most stable form, 1 atm, 25 °C
- solutions: 1 M concentration

Some common standard states:

H <sub>2</sub>	Fe	Hg
Cl <sub>2</sub>	Br <sub>2</sub>	I <sub>2</sub>
C(graphite)		

**Standard enthalpy of formation:  $\Delta H_f^\circ$**

enthalpy change when 1 mol of a compound is formed from its elements in their standard states


Write the chemical equation for  $\Delta H_f^\circ$  of NaCl(s)

Write the chemical equation for  $\Delta H_f^\circ$  of NH<sub>3</sub>(g)

**$\Delta H_f^\circ$  for any element in its standard state = 0**

Using standard enthalpies of formation

Calculate  $\Delta H^\circ$  for the following reaction using standard enthalpies of formation:



**Compound    $\Delta H_f^\circ$  (kJ/mol)**

NH <sub>3</sub> (g)	-45.9
O <sub>2</sub> (g)	
NO(g)	90.3
H <sub>2</sub> O(g)	-241.8

$$\Delta H_{\text{rxn}}^\circ = \sum n \Delta H_f^\circ (\text{products}) - \sum n \Delta H_f^\circ (\text{reactants})$$